



# Hepatoprotective Activity of 6-Heptadecylcyclohex-3-ene-1 Carboxylic Acid Isolated from the Methanol Extract of *Dichrostachys cinerea* Wight & Arn. Stem Bark

P. Suresh Babu <sup>a\*</sup>, V. Krishna <sup>b</sup> and D. C. Bhavya <sup>c</sup>

<sup>a</sup> Department of Phytochemistry and Pharmacology Laboratory, Trans-Disciplinary Research Foundation, Shankaraghatta – 577 451, Karnataka, India.

<sup>b</sup> P.G. Department of Studies and Research in Biotechnology and Bioinformatics, School of Biological Sciences, Kuvempu University, Shankaraghatta – 577 451, Karnataka, India.

<sup>c</sup> Department of P.G. Studies and Research of Microbiology, University of Davanagere, 577002, Karnataka, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/EJMP/2023/v34i21123

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94910>

Original Research Article

Received: 20/10/2022

Accepted: 24/12/2022

Published: 04/03/2023

## ABSTRACT

**Aim of the Study:** The phytoconstituent 6-heptadecylcyclohex 3-ene-1 carboxylic acid isolated from the methanol extract of *Dichrostachys cinerea* Wight. stem bark was evaluated for hepatoprotective activity against CCl<sub>4</sub> induced toxicity.

**Materials and Methods:** The constituent 6-heptadecylcyclohex 3-ene-1 carboxylic acid isolated from the methanolic extract of *D. cinerea* and the structure was confirmed by spectroscopic studies.

\*Corresponding author: E-mail: [spbioinfo@yahoo.co.in](mailto:spbioinfo@yahoo.co.in);

Hepatoprotective property was screened in male wistar strain rats. The parameters studied were estimation of liver function serum markers such as serum total bilirubin, total protein, alanine transaminase, aspartate transaminase, alkaline phosphatase and histological profile of the liver tissue.

**Results:** The LD<sub>50</sub> of methanolic extract and constituent, 6-Heptadecylcyclohex-3-ene-1 carboxylic acid were evaluated and found to be 500 and 100 mg/kg body weight respectively. The hepatoprotective activity of constituent was more significant as similar to the standard hepatoprotective drug silymarin. The histological profile of the liver tissue showed the presence of normal hepatic cords, absence of necrosis and fatty infiltration as similar to the controls.

**Conclusion:** The methanolic extract of *D. cinerea* stem bark and the phytoconstituent 6-heptadecylcyclohex-3-ene-1 carboxylic acid showed significant protection from CCl<sub>4</sub> induced liver damage.

**Keywords:** *Dichrostachys cineria*; hepatoprotective; 6-Heptadecylcyclohex-3-ene-1 carboxylic acid.

## 1. INTRODUCTION

Green plants possess the broad spectrum of synthetic activity and have been the source of many medically significant chemical compounds. *Dichrostachys cinerea* (Mimosae) is a deciduous medium sized tree commonly distributed in the forests of Africa, Australia, India and parts of South East Asia. It is commonly known as "sickle pod", "acacia Saint Domingue," "aroma," in English and "Vada" among the Kannada speaking people of Karnataka in southern India. "As per the traditional and folklore claims bark and the leaves of the *D. cinerea* were the potential source of drugs for ailments such as jaundice, inflammations rheumatism, fever, asthma, body ache, chest problems, toothache, ulcers, wounds eye diseases and aphrodisiac" [1]. "Preliminary phytochemical analysis of the plant specially leaves and bark extract contains flavonoids, tannins, triterpens, saponins and steroids" [2]. "The heartwood of *D. cinerea* contains aliphatics and triterpenoids" [3]. "Anti bacterial activity of the tannins isolated from the *D. cinerea*" [4]. In the present investigation 6-Heptadecylcyclohex-3-ene-1 carboxylic acid was isolated from the methanol extract of *D. cinerea* and its hepatoprotective property was screened against CCl<sub>4</sub> induced hepatic damage in rats.

## 2. MATERIALS AND METHODS

### 2.1 Plant Materials

The stem bark of *Dichrostachys cinerea* was collected from the Malebennur reserve forest range of Davanagere District, Karnataka, India. Taxonomic authenticity was confirmed by referring to herbarium specimen at Madras herbarium,

Botanical Survey of India, Southern Circle, Coimbatore and a voucher specimen (FDD-53) is deposited at Kuvempu University herbaria, Shankaraghatta Shivamogga Karnataka, India.

### 2.2 Isolation

The fresh stem bark of *D. cinerea* chapped, shade dried, powdered mechanically and was extracted using soxhlet apparatus with methanol for about 48 hrs. (200gm x 5). The extract was filtered and concentrated in vacuum under reduced pressure using rotary flash evaporator (Buchi, Flawil, Switzerland) and allowed it complete evaporation of the solvent. The yield was 2.16% (w/w).

Crude methanolic extract of stem bark was subjected for Thin Layer Chromatography (TLC) studies, which showed separation of single spot with the R<sub>f</sub> value 0.71. The active constituent was separated by column chromatography (180 g silica gel of 60 -120 mesh, 60x4 cm) and eluted with the solvent system toluene: ethyl acetate: diethyl ether in the ratio of 7:2:1. About 5 ml of collected fractions were monitored by TLC to check further separation and purity of separated constituent, which confirmed the single spot. The compound was re-crystallized in toluene and the yield was 230mg /5gms. The characterization of the compounds was done by IR, MASS, <sup>1</sup>HNMR and <sup>13</sup>C NMR spectroscopic studies.

### 2.3 Drug Formulations

Oral suspensions of the methanolic extract (50 mg/ml) and the isolated constituent 6-Heptadecylcyclohex-3-ene-1 carboxylic acid (10 mg/ml) was prepared in gum tragacanth (1 % w/v).

## 2.4 Animals

“Male Wistar albino rats weighing 150-200 g were procured from Central Animal House, National College of Pharmacy, Shivamogga and were maintained at standard housing condition. The animals were fed with commercial diet (Pranav Agro Industries Ltd., Sangli) and water ad- Libitum during the experiment. Acute toxicity study was conducted according to "staircase" method" [5]. “The LD<sub>50</sub> of methanolic extract and its constituent, 6-Heptadecylcyclohex -3-ene-1 carboxylic acid were found to be 500 and 100 mg/kg body weight respectively. One tenth of these doses (50 and 10 mg/kg, body weight respectively) was selected as the therapeutic dose for the evaluation of hepatoprotective activity”. [6]

## 2.5 Evaluation of Hepatoprotective Activity

“The animals were divided into 5 groups of 6 each. The animals of group I (control) received the vehicle gum tragacanth (1 ml/kg/day; 1 % w/v). Carbon tetrachloride with olive oil (1:1) was administered to all the animals of groups II to V in the dose of 0.1 ml/kg/day, ip for 14 days. The group III animals were treated with the standard drug silymarin (Ranbaxy Lab, Dewas;100 mg/kg/day, po). The animals of group IV received methanolic extract (50 mg/kg/day, po) and the animals of group V received the constituent 6-Heptadecylcyclohex -3-ene-1 carboxylic acid. (10mg/kg/day, po)” [7]. “The drugs were administered concomitantly for 14 days. The animals of all the groups were sacrificed on 14<sup>th</sup> day under light ether anesthesia. The blood sample of each animal was collected separately by carotid bleeding into sterilized dry centrifuge tubes and allowed to coagulate for 30 min at 37°C. The clear serum was separated by centrifugation at 2500 rpm for 10 min and subjected to biochemical investigation viz., total bilirubin” [7] total protein [8] serum alanine transaminase, aspartate transaminase,[9], and alkaline phosphatase [10] Results of biochemical estimations were expressed as mean±SE of six animals in each group. The statistical analysis was carried out using one way ANOVA. The difference in values at P≤ 0.05 was considered as statistically significant.

## 2.6 Study of Histopathology

“The liver samples were excised from the animals of each group after draining the blood

and washed with the normal saline. Initially the materials were fixed in 10% buffered neutral formalin for 48 hrs. They were processed for paraffin embedding. The sections were taken at 5 μm thickness, processed in alcohol-xylene series and were stained with alum-haematoxylin and eosin” [11,12]. The sections were examined microscopically for the evaluation of histological changes.

## 3. RESULTS

The compound isolated from stem bark of methanolic extract of *D .cinerea* is a light yellow amorphous powder; Melting point is recorded to be 76 -79 °C.

The IR spectrum exhibits bands at 3341 cm<sup>-1</sup> for -OH group,1736 cm<sup>-1</sup> for carboxylic group, 1647 for C=C (Fig. 1).

The molecular formula of the compound is C<sub>25</sub>H<sub>46</sub>O<sub>2</sub> evident from its mass spectrum by exhibiting a pseudomolecular ion [M-H]<sup>-</sup> at m/z 376.90. The peaks were observed at m/z355,m/z184 m/z170and m/z130(Fig. 2).

The <sup>1</sup>H NMR spectra (CDCl<sub>3</sub> +DMSO.D6 MIX.) showed peaks at 5.06(1H,s,OH), 2.0(14H,,J=6.8Hz, CH<sub>2</sub>)0.66(3H,t,J=6.80 Hz,CH<sub>3</sub>)0.88[28H,s,14xCH<sub>2</sub>]. The peaks at δ 0.67 (ppm) for terminal methyl and δ 1.67 (ppm) for a methyl group attached to a double bond. The signal at δ 2.60 is due to a proton attached to a carboxyl group in a ring system. The multiplet at δ 5.54 is due to the presence of an unsaturated proton. In addition, a strong singlet represents the long chain methylene groups (Fig. 3).

The <sup>13</sup>C-NMR spectra exhibits two signals at δ 127.60 and δ 129.81 were due to the presence of a double bond. The singlet at δ 55.70 and δ 55.10 are due to carbon atoms of long chain methylene groups. The other signals at δ 13.83, δ 22.32, δ 24.69, δ 26.84, δ 28.86, are due to long chain alkyl group (Fig. 4).

Based on above facts the structure of the compound written as shown in the Fig. 5 and the compound was identified as 6-Heptadecylcyclohex -3-ene-1 carboxylic acid.

After 14 days treatment, biochemical analysis of blood samples of CCl<sub>4</sub> treated animals showed significant increase in the levels of total bilirubin (6.22 fold), aspartate transaminase (5.71 fold), alanine transaminase(4.22 fold) and alkaline

phosphatase (3.47 fold) as compared to the controls. In addition, the total protein level (44.86%) was decreased reflecting the liver injury due to the toxic effect of CCl<sub>4</sub>. The blood samples of the animals treated with the methanolic extract and the constituent 6-Heptadecylcyclohex -3-ene-1 carboxylic acid

were showed significant reduction in the levels of liver function serum markers. The effect was more pronounced in the animals treated with 6-Heptadecylcyclohex -3-ene-1 carboxylic acid as similar to the effect of the standard drug silymarin (Table 1).




Fig. 1. IR Spectrum– 6-Heptadecylcyclohex–3-ene-1 carboxylic acid




Fig. 2. MR Spectrum of 6-Heptadecylcyclohex–3-ene-1 carboxylic acid

Table 1. Effect of methanolic extract and its constituent 6-heptadecylcyclohex 3-ene-1 carboxylic acid on CCl<sub>4</sub> induced hepatotoxicity

| Groups                                       | T. BIL (mg/dl) | Total protein | AST (IU/L)    | ALT (IU/L)   | ALP (IU/L)   |
|--|----------------|---------------|---------------|--------------|--------------|
| Control                                      | 0.36±0.08      | 7.40±0.36     | 209.41±13.57  | 76.31±9.11   | 116.86±9.57  |
| CCl <sub>4</sub>                             | 2.24±0.29      | 4.08±0.27     | 1194.08±73.60 | 321.67±27.93 | 403.73±8.66  |
| Silymarin+ CCl <sub>4</sub>                  | 0.40±0.14      | 7.34±0.42     | 217.64±23.8   | 78.81±8.05   | 118.21±9.72  |
| Methanol extract + CCl <sub>4</sub>          | 0.58±0.14      | 6.97±0.38     | 310.14±36.96  | 106.50±8.08  | 144.60±18.68 |
| 6-heptadecylcyclohex 3-ene-1 carboxylic acid | 0.46±0.12      | 7.18±0.17     | 231.72±22.07  | 82.95±7.17   | 120.01±9.16  |
| F value                                      | 22.4           | 18.1          | 114           | 53.8         | 113          |

Values are mean ± SE; n=6 in each group; P<0.05 when compared to control




Fig. 3. <sup>1</sup>H NMR of 6-Heptadecylcyclohex-3-ene-1 carboxylic acid




Fig. 4. <sup>13</sup>C NMR of 6-Heptadecylcyclohex-3-ene-1 carboxylic acid




Fig. 5. Structure of 6-Heptadecylcyclohex-3-ene-1 carboxylic acid

The histological profile of control animal showed normal hepatocytes (Fig. 6A). The section of liver of the animals treated with CCl<sub>4</sub> exhibited intense centrilobular necrosis, vacuolization and macro vesicular fatty changes (Fig. 6B). The liver sections of silymarin treated animals showed normal hepatic architecture (Fig. 7A). Moderate accumulation of fatty lobules was observed in the

liver sections (Fig. 7B) of methanolic extract treated animals. The liver sections of the animals treated with the constituent 6-Heptadecylcyclohex -3-ene-1 carboxylic acid exhibited significant liver protection against CCl<sub>4</sub> induced liver damage as evident by the presence of normal hepatic cords, absence of necrosis and fatty infiltration (Fig. 7C).




Fig. 6.




Fig. 7.

Figs. 6-7. Histological slides

#### 4. DISCUSSION

Medicinal plants are an integral component of research developments in the pharmaceutical industry. Furthermore, an increasing use of medicinal plants in the pharmaceutical industries

have been traced to the extraction and development of several drugs from plants as well as from traditionally used rural herbal remedies. The constituent 6-Heptadecylcyclohex -3-ene-1 carboxylic acid isolated from the stem bark was showed significant hepatoprotective activity

against  $\text{CCl}_4$  induced hepatotoxicity. To the best of our knowledge the 6-Heptadecylcyclohex-3-ene-1 carboxylic acid is a novel phytoconstituent isolated for the first time from *D. cinerea*. Similar fatty acid derivatives such as 5-hexyl-2-(8-oxooctyl)cyclohexane 1-carboxylic acid (Pubchem CID23345599) with MW: 354.524020 g/mol | MF:  $\text{C}_{21}\text{H}_{38}\text{O}_4$  3-Heptylcyclohexane-1-carboxylic acid (PubchemCID 20320740) with MW 226 355040 g/mol | MF:  $\text{C}_{14}\text{H}_{26}\text{O}_2$  . 2- butylcyclohexane-1-carboxylic acid (PubchemSID53962009) has MW: 184g/mol g/mol |MF: $\text{C}_{11}\text{H}_{20}\text{O}_2$  and 2-butylcyclohexane-1-carboxylic acid MW: 184.275300 g/mol | MF:  $\text{C}_{11}\text{H}_{20}\text{O}_2$  (PubchemCID 12933617) have been reported. Reports also indicated that  $\omega$ -3 and  $\omega$ -6 fatty acids exhibited significant hepatoprotective property in rats [13] and supplementation of omega-3 fatty acid would reduce hepatocellular damage and cell death in a model of murine common bile duct ligation [14,15].

It is well established that  $\text{CCl}_4$  induces hepatotoxicity by metabolic activation, therefore it selectively causes toxicity in liver cells maintaining semi-normal metabolic function [16,17].  $\text{CCl}_4$  is bio-transformed by the cytochrome P450 system in the endoplasmic reticulum to produce trichloromethyl free radical ( $\cdot\text{CCl}_3$ ). Trichloromethyl free radical when combined with cellular lipids and proteins in the presence of oxygen form trichloromethyl peroxy radical, which may degrade lipids on the membrane of endoplasmic reticulum, thus leads to elicit lipidperoxidation and finally results in cell death [14]. In this present study it was noted that the administration of  $\text{CCl}_4$  decreased the levels of total protein and increased levels of marker enzymes. The estimation of total bilirubin depicts the depth of jaundice and the elevated level of marker enzymes, ALT, AST and ALP indicates the intensity of liver damage. Marked elevation in the levels of total bilirubin and marker enzymes in the serum of  $\text{CCl}_4$  intoxicated rats (1194.08 $\pm$ 73.60, 321.67 $\pm$ 27.93 and 403.73 $\pm$ 8.66 respectively) were significantly decreased in the serum of the animals treated with methanol extract (310.14 $\pm$ 36.96, 106.50 $\pm$ 8.08 and 144.60 $\pm$ 18.68 respectively) and 6-Heptadecylcyclohex-3-ene-1 carboxylic acid (231.72 $\pm$ 22.07, 82.95 $\pm$ 7.17 and 120.01 $\pm$ 9.16 respectively). "The protection against the injurious effects of carbon tetrachloride that may result from the interference with cytochrome P450, resulting in the hindrance of the formation of hepatotoxic free radicals. The site-specific oxidative damage in some susceptible amino

acids of proteins may regarded as the major cause of metabolic dysfunction of liver during pathogenesis" [18]. "Bilirubin is the conventional indicator of liver diseases" [19]. "The restoration bilirubin levels may be due to the inhibitory effects on cytochrome P450 or/and promotion of its glucuronidation" [20]. The attainment of these marker enzymes towards a near-normalcy in the animals treated with methanol extract and its isolated phytoconstituent 6-heptadecylcyclohex-3-ene-1 carboxylic acid confirms the hepatoprotective effect. The results were found comparable to a commercial hepatoprotective drug silymarin which is a composite name of three flavonoids isolated from milk thistle *Silybum maritimum*. Among all the treated groups the hepatoprotective activity was more in the animals treated with phytoconstituent 6-heptadecylcyclohex-3-ene-1 carboxylic acid. Similar type of investigations on the effect of phytoconstituents isolated from the medicinal plants in restoring the levels of liver function markers was reported [21-24].

## 5. CONCLUSION

In conclusion, the constituent 6-heptadecylcyclohex-3-ene-1 carboxylic acid isolated from the methanolic extract of *D. cinerea* is a novel phytoconstituent afforded protection from  $\text{CCl}_4$  induced liver damage. The possible mechanism that it may protect the liver by  $\text{CCl}_4$  toxicity or may act as a free radical scavenger intercepting those radicals involved in  $\text{CCl}_4$  metabolism by microsomal enzymes. By trapping oxygen related free radicals the compound could hinder their interaction with polyunsaturated fatty acids and would abolish the enhancement of lipid peroxidative processes. The present investigation also supports the ethnomedical uses of *D. cinerea*.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

The Institutional Animal Ethical Committee (Reg.No.144/NCP/IAEC/CLEAR/P.COL.3/2006-07) permitted the study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Bako SP, Bakfur MJ, John I, Bala EI. Ethnomedical and phytochemical profile of some savanna plant species in Nigeria. *International Journal of Botany*. 2005; 1(2):147-150.
- Igoli JO, Igwue IC, Igoli NP. Traditional medicinal practices among the Igede people of Nigeria. *Journal of Herbs, Spices & Medicinal Plants*. 2004;10(4):1049.
- Jain R, Saxena U. Aliphatics and triterpenoid's from the heartwood of *Dichrostachys cinerea*. *Journal of the Indian Chemical Society*. 2003;80(6):656-658.
- Banso A, Adeyemo SO. Evaluation of antibacterial properties of tannins isolated from *Dichrostachys cinerea*. *African Journal of Biotechnology*. 2007;6(15): 1785-1787.
- Ghosh MN. *Fundamentals of experimental pharmacology*, 2nd ed. Scientific Book Agency Kolkatta. 1984;154.
- Meharie BG, Amare GG, Belayneh YM. Evaluation of hepatoprotective activity of the crude extract and solvent fractions of *Clusia abyssinica* (Euphorbiaceae) Leaf against CCl<sub>4</sub>-Induced hepatotoxicity in Mice. *J Exp Pharmacol*. 2020;12:137-150. Available:<https://doi.org/10.2147/JEP.S248677>
- Mallory HT, Evelyn EA. The determination of bilirubin with photoelectric colorimeter. *Journal of Biological Chemistry*. 1937;119:481-490.
- Kingsley SR, Frankel SJ. The determination of serum total protein albumin and globulin by the biuret reaction. *Journal of Biological Chemistry*. 1939;128:131-37.
- Reitman S, Frankel S. A Colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminase. *American Journal of Clinical Pathology*. 1957;28:53-63.
- Bessey OA, Lowery DH, Brock MJ. A method for the rapid determination of alkaline phosphatase with five cubic meters of serum. *Journal of Biological Chemistry*. 1964;164:321-29.
- De S, Suresh R, Babu AMSS, Aneela S. *in-vivo* hepatoprotective activity of methanolic extracts of *Sphaeranthus amaranthoides* and *Oldenlandia umbellata*. *J Pharmacogn*. 2017;9(1).
- Galigher AE, Kayloff EN. Essentials of practical microtechniques. *Journal of Pharmacological Research*. 1997;36(4): 315-321.
- Guntupalli M, Chandana V, Pushpangadan P, Shirwaiker AI. Hepatoprotective effects of rubiadin, a major constituent of *Rubia cordifolia* Linn. J. *Ethnopharmacol*. 2006;103:484–490. [PubMed] [Google Scholar]
- Makni M, Fetoui H, Gargouri NK, Garoui EM, Jaber H, Makni J, Boudawaraand T, Zeghal N. Hypolipidemic and hepatoprotective effects of flax and pumpkin seed mixture rich in ω-3 and ω-6 fatty acids in hypercholesterolemic rats. *Food and Chemical Toxicology* 2008;46(12):3714-3720.
- Sang Lee, Sendia Kim, Hau D. Le, Jonathan Meisel, Robbert A.M. Strijbosch, Vania Nose, Mark Puder. Reduction of hepatocellular injury after common bile duct ligation using omega-3 fatty acids. *Journal of Pediatric Surgery*. 2008;43:2010-2015.
- Mujumddar AK, Upadhye AS, Pradhan AM. Effect of *Azadirachta indica* leaf extract on CCL<sub>4</sub> Induced hepatic damage in albino rats. *Indian Journal of Pharmacological Science*. 1998;60:363.
- Opoku AR, Ndlovu IM, Terblanche SE, Hutchings AH. *in vivo* hepatoprotective effects of *Rhoicissus tridentata* sub sp. *cuneifolia*, a traditional Zulu medicinal plant, against CCl<sub>4</sub>-induced acute liver injury in rats. *South African Journal of Botany*. 2007;73:372–377.
- Uday bandyopadhyay, Dipak D, Banerji ranjit K. Reactive oxygen species: Oxidative damage and pathogenesis. *Current Science*. 1999;5:658.
- Barnes S, Gallo GA, Trash DB, Morris JS. Diagnostic value of serum bile acid estimations in liver disease. The BMJ Publishing Group Ltd & Association of Clinical Pathologists. *Journal of Clinical Pathology*. 1975;28:506-509.
- Cavin C, Mace K, Offord EA, Schilter B. Protective effects of *Coffee diterpenes* against aflatoxin B1-induced genotoxicity: Mechanisms in rat and human cells. *Food Chemistry and Toxicology*. 2001;39:549-556.
- Saraswath B, Visen PKS, Patnaik GK, Dhawan BN. Anticholestic effect of picroliv, active hepatoprotective principle of *Picrorhiza kurrooa*, against carbon

- tetrachloride induced cholestatis. Indian Journal of Experimental Biology. 1993; 31:316.
22. Du XM, Sun NY, Hayashi J, Chen Y, Sugiura M, Shoyama Y. Hepatoprotective and antihyperliposis activities of *in vitro* cultured *Anoectochilus formosanus*. Phytotherapy Research. 2003;17:30–33.
23. Singh B, Chandan BK, Prabhakar A, Taneja SC, Singh J, Qazi GN. Chemistry and hepatoprotective activity of an active fraction from *Barleria prionitis* Linn. in experimental animals. Phytotherapy Research. 2005;391.
24. Das S, Sarma G, Barman S. Hepatoprotective activity of aqueous extract of fruit pulp of *Cassia fistula* (AFCF) against Carbon Tetrachloride (CCL<sub>4</sub>). Journal of Clinical and Diagnostic Research. 2008;2:1133-1138.

© 2023 Babu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/94910>